



ACNP

American College of Neuropsychopharmacology

EMBARGOED FOR RELEASE

December 9, 2014 at 12:01 a.m. Eastern

CONTACT: ACNP, (615) 324-2360

acnp@acnp.org

Sleep Disturbance Linked to Amyloid in Brain Areas Affected by Alzheimer's Disease

Phoenix, AZ (December 9th, 2014) - Healthy, elderly research participants who report being more sleepy and less rested have higher levels of amyloid deposition in regions of the brain that are affected in Alzheimer's disease, according to a report presented today at the American College of Neuropsychopharmacology annual meeting in Phoenix (Arizona). If sleep disturbance is a cause of amyloid accumulation, it may be an early target for intervention to prevent the progression of cognitive deficits in late life.

Numerous studies have shown the importance of sleep and the effect that sleep deprivation can have on our brains. Sleep apnea is associated with cognitive dysfunction across the lifespan and both untreated apnea and sleep disturbance are associated with increased cognitive dysfunction in patients with Alzheimer's disease. Evaluating physical changes in the brain and how they are related to sleep quality can help determine the extent to which sleep can provide a window on brain function and pathology.

Using positron emission tomography (PET) scanning with a tracer that visualizes deposits of amyloid, a protein that is elevated in the brains of Alzheimer's disease patients, Ruth Benca and her colleagues at the University of Wisconsin-Madison studied the relationship between sleep quality and brain amyloid levels in a group of 98 cognitively healthy volunteers, 50-73 years of age. The subjects, who were participants in the Wisconsin Registry for Alzheimer's Prevention, completed questionnaires about their sleep and problems related to it. Those who reported greater sleepiness showed greater amyloid accumulation in areas of the cerebral cortex that are heavily affected in Alzheimer's disease — the supramarginal and frontal medial orbital areas. Higher amyloid in these regions was also linked to less restful sleep and more sleep problems.

Although it is tantalizing to speculate that the link between sleep disturbances and amyloid deposition in the brain may help us identify an early, modifiable marker for Alzheimer's disease, it is too soon to draw that conclusion. "We still need to determine whether sleep disturbance promotes amyloid deposition in the brain, or if a neurodegenerative process produces disordered sleep", commented Dr. Benca. Future work is needed to answer that question and to determine whether interventions that improve sleep can prevent brain changes that lead to Alzheimer's disease.

This research was supported by funding from the National Institutes of Health: National Institute on Aging, National Institute of General Medical Sciences, and the Clinical and Translational Science Award Program.

###

ACNP, founded in 1961, is a professional organization of more than 700 leading scientists, including four Nobel Laureates. The mission of ACNP is to further research and education in neuropsychopharmacology and related fields in the following ways: promoting the interaction of a broad range of scientific disciplines of brain and behavior in order to advance the understanding of prevention and treatment of disease of the nervous system including psychiatric, neurological, behavioral and addictive disorders; encouraging scientists to enter research careers in fields related to these disorders and their treatment; and ensuring the dissemination of relevant scientific advances.